

lemmas play a crucial role.

Lemma 1. *Let g be an element of G and let $\{W_1, W_2, \dots, W_m\}$ be a covering of G by evenly covered neighborhoods such that $g \in W_1 \setminus \bigcup_{k=2}^m W_k$. Let $p^{-1}(W_1) = \bigcup_{l=1}^n V_l$ and $p^{-1}(g) \cap V_1 = \{y_0\}$. Then there exists a neighborhood $V_0 \subset V_1$ of the point y_0 satisfying the following property: if $T_{t_0}(V_0) \cap V_0 \neq \emptyset$ for some $t_0 \in R$, then $T'_{t_0}(V_0) \subset V_1$.*

Lemma 2. *An orbit $R_x = \{T_t(x) : t \in R\}$ of each point $x \in X$ is dense in the connected space X .*

Finally, we formulate the theorem on covering groups for compact solenoids.

Theorem. *Let $p : X \rightarrow G$ be an n -fold covering of compact solenoidal group G by a connected topological space X . Then there exists a group structure in X turning $p : X \rightarrow G$ into a homomorphism between compact abelian groups.*

The work is supported by RFFI (project 99-01-00441) and NIOKR of Ta- tarstan.

REFERENCES

1. Grigorian S. A. *Generalized analytic functions*, (in Russian)// Uspekhi Matem. Nauk. – 1996. – V. 49. – No 2. – P. 3–42.
2. Grigorian S. A., Gumerov R. N. and Kazantsev A. V. *Group structure in finite coverings of compact solenoidal groups*, // Lobachevskii J. of Math., to appear.
3. Pontryagin L. S. *Continuous groups* (in Russian). – Moscow: Nauka, 1984.

A. R. Kacimov (Muskat, Oman)

EXTREME PROPERTIES OF THE TAYLOR-SAFFMAN CURVE

Since the seminal Taylor-Saffman experiments and theoretical

investigations of viscous fingers in the Hele-Shaw apparatus many attempts have been done to explain the mechanisms of so-called "selection". Still, it is unclear why practically a half-channel finger appears though mathematically a whole family of interfaces is possible. We apply the method of inverse boundary-value problems [1] and arrive at the Taylor-Saffman shapes from solution of optimal shape design problems [2,3] illustrating that "selected" curves possess fascinating extreme properties. In particular, we show that:

a) The Taylor-Saffman bubble is a limiting case of the Taylor-Saffman finger.

b) The Taylor-Saffman bubble coincides with the Polubarinova-Kochina subsurface contour of a concrete dam of constant hydraulic gradient, which is a solution of an isoperimetric problem on maximum of the cross-sectional area of the dam.

c) The Taylor-Saffman half-channel finger coincides with the Morse-Feshbach [4] "extreme" boundary of a variable condenser. Hence, the finger is an equipotential line generated by a semi-infinite linear source.

d) The Taylor-Saffman finger coincides with an abrupt interface between seeping fresh and stagnant saline water in a polder-type system [5], i.e., the finger is a stream line of a vortex generated flow.

Obtained explicit analytic solutions of optimization problems are also used for estimations of integral and local flow characteristics (total flow rate and field intensity).

REFERENCES

1. Ilyinsky N. B., Kacimov A. R. *The estimation of integral seepage characteristics of hydraulic structures in terms of the theory of inverse boundary-value problems*// Zeitschr. angew. Math. Mech. – 1992. – B. 72. – No 2. – P. 103–112.
2. Kacimov A. R. *Optimization of the protrusion shape for a Couette type flow*// Optimal Control Applications and Methods. – 1994. – V. 15. – P. 193–203.
3. Kacimov A. R. *Optimal shape of a variable condenser*// Proc. Royal Soc. London A, in press.
4. Morse P. M. and Feshbach H. *Methods of Theoretical Physics*. V. 2. – New York: McGraw-Hill, 1953.

5. Strack, J.D.L. *Groundwater Mechanics*. Prentice Hall, Englewood Cliffs, 1989.

I. I. Sakhaev (Kazan), C. Lomp (Porto),
M. F. Nasrutdinov (Kasan)

ON PROJECTIVE MODULES OF FINITE DUAL GOLDIE DIMENSION

Let R be an associative ring with unit, P will be a left unital R -module and $J(P)$ denote the Jacobson's radical of P .

A submodule N of P is said *small* in P if for every submodule U of P the equation $N + U = P$ involves $U = P$. A module P is said to be hollow if $P \neq 0$ and every proper submodule of P is small in P . A module P is said to have *finite hollow dimension* (or *finite dual Goldie dimension*) if there exists an exact sequence

$$P \xrightarrow{g} \bigoplus_{i=1}^n H_i \longrightarrow 0$$

where all H_i are hollow and the kernel of R -homomorphism g is small in P . Then n is called the hollow dimension of P and we write $hdim(P) = n$.

In the paper [1] it was formulated the question: *Is every projective R -module P with semilocal endomorphism ring $Hom_R(P, P)$ finitely generated?*

If P is projective R -module and $Hom_R(P, P)$ is semilocal ring then by the theorem 3.10 [1] we have $hdim_R(P) < \infty$. We have proved:

Theorem *Let P be a projective R -module and $Hom_R(P, P)$ is semilocal ring. Then the following conditions are equivalent:*

- (a) P is finitely generated.
- (b) $hdim_R(P) = hdim_R(P/J(P))$.

Research of I.I. Sakhaev and M.F. Nasrutdinov is supported by RFFI grant 99-01-00469.